

Immediate effects of longitudinal vs. Transverse tibial nerve massage on vibration perception thresholds and thermal perception thresholds in asymptomatic subjects: A pilot randomized clinical trial (UTRN 014427448-2101200878203)

Kumar P. S.*

Adhikari P.**

Jeganathan P.S.***

ABSTRACT

Purpose of study: To compare the immediate effects of two techniques of nerve massage- longitudinal and transverse, for tibial nerve on the vibration and temperature thresholds in the foot in asymptomatic subjects. **Materials and methods:** 48 subjects of either sex (34 male, 14 female), with mean age 23.6 ± 4.8 years, volunteered to participate by incidental sampling after giving informed consent. The side of the leg for first intervention was chosen by simple random sampling and the choice of first intervention (longitudinal or transverse nerve massage) was allocated in a concealed manner using block randomization. One intervention was a longitudinal nerve massage (to either side) while other was a transverse nerve massage (to the other side). One tester administered the nerve massage techniques while an independent blinded observer measured the outcome measures. Both the interventions lasted for 2 minutes each. The techniques were applied for 10 reps and 2 sets. Pre, immediate post intervention and post intervention- 5 mins assessment of vibration thresholds, heat thresholds and cold thresholds of the sole of the foot were assessed using a Vibrotherm® (Biothesiometer) by the blinded observer.

Results: Tibial nerve transverse massage technique reduced the vibration thresholds ($2.1 \pm .84$ volts) in the foot significantly better ($p = .000$) than that of the longitudinal nerve massage. Transverse massage technique also reduced the heat thresholds- 6.4 ± 1.8 degrees Celsius ($p = .054$) and increased the cold thresholds- 4.8 ± 2.2 degrees Celsius ($p = .87$) clinically but not statistically significant ($p > .05$). Both effects were however reversible within five minutes ($p < .05$). There was no effect of side of dominance ($p = .14$) in all the subjects studied. **Conclusion:** Transverse nerve massage technique was significantly better than the longitudinal massage technique in reducing the vibration thresholds for tibial nerve. Clinically, transverse nerve massage technique also reduced the heat thresholds and increased the cold thresholds for tibial nerve in the subjects studied.

Keywords: Neurodynamics, nerve massage, quantitative sensory testing, randomized controlled trial

INTRODUCTION

Nervous system and its role in pain production has been increasingly identified and addressed in musculoskeletal physical therapy practice using

neurodynamic assessment and treatment methods which focus on nerve mobility- both intraneural (between neural connective tissue sheaths) and extraneural (between nerve and its surrounding structures).¹

Symptoms arising from peripheral nerves have been categorized into positive symptoms (dysesthetic pain such as hyperalgesia, allodynia, tingling, numbness, paresthesia and/or shooting pain) and negative symptoms (with neurological deficits such as sensory loss, motor loss and reflex loss) and appropriate management was indicated to suit individual cases with clinical reasoning.²

Author's Affiliations: *Associate professor in Musculoskeletal and Manual Therapy, Dept of Physiotherapy, **Professor and Unit-I Head, Dept. of Medicine, ***Professor, Dept of Physiology, Kasturba Medical College (Manipal University), Mangalore.

Reprint requests: P. Senthil Kumar, (PhD) P.T., Associate professor in Musculoskeletal and Manual Therapy, Dept of Physiotherapy, Kasturba Medical College (Manipal University), Mangalore. senthil.kumar@manipal.edu

(Received on 22.04.10, accepted on 19.05.10)

Neurodynamic method encompasses manual techniques which address and/or treat nerve mobility (both intraneural and extraneural) in both directions- longitudinal and transverse.³ Traditionally neurodynamics, as it was developed consisted of techniques which focused on longitudinal nerve mobility testing and treatment, and hence termed as “neural tension tests”.⁴ Much later the emphasis shifted course to transverse mobility, which was detected by manual palpation of the nerve along its course, owed its contribution to development of “nerve massage” techniques.⁴

Over the course of clinical manual therapy practice, newer techniques have been developed modifying the existing neurodynamic techniques when performed in different combinations and in different sequence of moving the joints crossed by the nerve. One technique which thrives on the neural movement along its interface structures without an increase in its “tension” was termed as “slider” techniques. The slider techniques improve gliding without compromising the nerve circulation or increasing nerve tension.⁵

Sliders and tensioners are two techniques which form an integral part of comprehensive neurodynamic assessment and treatment in manual therapy which addresses symptoms originating from peripheral nerves. Biomechanically, the slider technique stretches the nerve at one end while relaxing at another; the tensioner technique stretches the nerve at both the ends and/or joints.⁶

Soft tissue mobilization and massage had been shown to improve tissue flexibility and mobility by increasing nutrition and relieving adhesions. Whilst various techniques of massage like kneading and effleurage was extensively studied using application on muscles, tendons and ligaments, deep transverse friction massage as advised by Cyriax,⁷ was being increasingly used for a variety of soft tissue conditions.

Neuromechanics^{8,9,10} and neurophysiology were two interdependent and interrelated phenomena which formed the basis for neurodynamic assessment and treatment methods. There were numerous studies on neurodynamic assessments including normal responses,^{11,12,13,14,15,16,17}

movements’ combinations for neurodynamic tests,^{18,19,20,21,22,23} in symptomatic subjects,^{24,25,26,27,28,29,30,31,32,33,34,35,36} and also as comparison between normals and symptomatic subjects.^{37,38} Other studies on neurodynamic treatment effects were on nerve gliding exercises for carpal tunnel syndrome,^{39,40,41,42,43,44,45,46,47,48,49} cubital tunnel syndrome,^{50,51} radial tunnel syndrome,⁵² lateral epicondylitis,^{53,54} thoracic outlet syndrome,^{55,56} cervical cord compression,⁵⁷ cervical radiculopathy,⁵⁸ cervicobrachial pain syndrome,^{59,60,61,62} non radicular low back pain,⁶³ lumbar nerve root irritation,⁶⁴ lower extremity symptoms⁶⁵ and lumbar spine surgery.⁶⁶

The concept of nerve and its mobility (both intraneural and extraneural), which assumes nerves as similar to that of any other soft tissues, proposed neural tissue metabolism much similar to that of any other tissue. Reduced mobility had the tendency to lead to accumulation of extracellular fluid, metabolites, and other tissue end products which need to be removed and circulated for proper nerve function. Massage techniques directed at other tissues might also have had their effects on the nerves indirectly which was less documented and/or less studied in the literature.

Statement of problem

Studies comparing effects of nerve massage in lower limb could not be retrieved from the existing literature. The comparative effects of the two techniques of nerve massage (longitudinal and/or transverse nerve massage) on vibration and temperature thresholds were also not documented in literature.

Purpose of study

To compare the effects of one session application of two nerve massage techniques- longitudinal and transverse massage for tibial nerve on vibration and temperature (heat and cold) thresholds in asymptomatic subjects.

Hypothesis of study

The effects of the two nerve massage techniques on vibration and temperature thresholds would be different in asymptomatic subjects.

MATERIALS AND METHODS

Study Design

Randomized Controlled Trial, with observer and tester blinded.

Study Setting

Health Lounge, Dept of Medicine, Kasturba Medical College Hospital, Attavar, Mangalore.

Study Duration

Four months, from August to November 2007.

Sampling

Volunteers were recruited on incidental sampling.

Sample size

48 subjects of either sex (34 male, 14 female), with mean age 23.6 ± 4.8 years, volunteered to participate by incidental sampling after giving informed consent.

Subjects

Volunteers were described about the conduct of the study and its significance for assessment and treatment implications in patients with lower extremity complaints.

Inclusion criteria

Subjects were deemed eligible if they did not have any symptoms in lower limb or low back region for the past 6 months (from day of assessment by tester).

Exclusion criteria

Excluded subjects consisted of inability of subjects to understand and/or co-operate during quantitative sensory testing and/or manual nerve mobility testing (on palpation) and/or treatment.

Tester

The primary investigator was an Assistant professor in Musculoskeletal and Manual Therapy, Dept of Physiotherapy, Kasturba Medical College, Mangalore, and Co-investigator was a Professor of Medicine, Kasturba Medical College, Mangalore. Independent blinded observer

was an intern in physiotherapy who was trained by the primary investigator for assessment and recording of quantitative sensory testing findings.

Testing Instruments


Vibrotherm™ Biothesiometer (Diabetic Foot Kare India, Chennai).

The assessment of vibration thresholds and temperature thresholds were done based on the instructions provided in the manual. After choosing the setting, the transducer probe is placed at one of the three sites in the sole of the foot (in a randomly selected order) and then instructed to report when he/she starts feeling the onset of sensation of vibration/ heat/ cold. The intensity knob was adjusted gradually, noting for the subject's response throughout the testing procedure. The point when the subject felt the onset of sensation, the reading on the digital scale of the biothesiometer would indicate the value of sensory threshold for that tested sensation. Three values were thus obtained which were then averaged to get a single value for that side for that sensation.⁶⁸ Biothesiometer assessment of quantitative sensory testing was validated highly and studied extensively for its high reliability and responsiveness.⁶⁹ Quantitative sensory testing was even proven to be better than nerve conduction studies in diagnosis and prognosis in peripheral nerve disorders.⁷⁰ Pilot intra and inter tester reliability was studied for the blinded observer for the biothesiometer prior to the actual study.

PROCEDURE

The side of the subject's leg was first chosen by simple random sampling (lots method) and the choice of first intervention (longitudinal or transverse nerve massage) was allocated in a concealed manner using block randomization (blocks of two for every 10 subjects). The subject's leg was administered either of the two techniques first, followed by the other technique next on the other leg.

Fig 1: CONSORT Flow Chart showing the flow of participants through the study



Tibial nerve longitudinal nerve massage technique:

The therapist stood at the side to be treated with the subject lying supine on a plinth. The therapist first manually palpates along the course of the tibial nerve both distoproximally and proximodistally for identifying the subcutaneous course of the tibial nerve from the popliteal fossa to the foot. Then the therapist applied distal to proximal directed force on the tibial nerve in short segments starting from upper leg and progressing to middle and lower leg segments. The hand

position of the therapist was that the proximal hand localized the nerve (by manual palpation) while the distal hand applied the force for massage towards the proximal hand. The massage was applied with the finger pads of 2nd, 3rd and 4th digits of the distal hand with the wrist slightly shifting from ulnar deviation to neutral during the technique. Two sets of ten repetitions of the technique was done at each segment. Total of six sets with each consisting of ten repetitions was thus applied to that side, with the intervention lasting 4 mins.



Tibial nerve transverse nerve massage technique:

The initial preparatory procedure was the same as that for the longitudinal nerve massage technique. Transverse nerve massage technique was applied with the finger tips of the 2nd, 3rd and



4th digits of the therapist's distal hand, while his wrist was shifting from flexion to extension during the technique application. Similar dosage was applied for this technique thus achieving equal treatment duration of 4 mins.



DATA COLLECTION

One tester administered the neurodynamic techniques while an independent blinded observer measured the three outcome measures. Pre, immediate post intervention and post intervention- 5 mins assessment of vibration thresholds, heat thresholds and cold thresholds of the sole of the foot were assessed using a Vibrotherm® (Biothesiometer) by the blinded observer. Only one trial was taken for each of the sites.

DATA ANALYSIS

The between side (intervention group- slider vs. tensioner) analysis of three outcome measures (vibration threshold, heat threshold, cold threshold) at three levels of measurement (pre, immediate post, post 5 min intervention) were analyzed using repeated measures ANOVA at 95% confidence interval using SPSS 12.0.1 for Windows.

DISCUSSION

Our study found that responses were different between the sides of lower extremities in the three


outcomes (vibration threshold, heat threshold and cold threshold) which was attributable to the intervention and not due to the influence of lower limb dominance ($p > .05$). The study reported better

RESULTS


	Tibial nerve Transverse Massage intervention Mean \pm SD			Tibial nerve Longitudinal Massage intervention Mean \pm SD			P value
	Pre	Post	Post 5 min	Pre	Post	Post 5 min	
Vibration threshold (Volts)	4.33 \pm 2.1	2.66 \pm 1.2	4.12 \pm 1.8	4.08 \pm 1.9	3.78 \pm 2.1	4.32 \pm 1.9	.000 ^{**}
Heat threshold (Degree Celsius)	46.4 \pm 8.8	44.4 \pm 14.8	47.7 \pm 8.2	44.7 \pm 10.3	43.9 \pm 13.3	45.5 \pm 11.1	.054
Cold threshold (Degree Celsius)	28.6 \pm 11.7	30.9 \pm 18.8	29.2 \pm 9.6	24.4 \pm 12.9	28.7 \pm 8.3	27.1 \pm 7.4	.087

^{**}- significant at $p < .05$ level.


Graph 1 showing pre, post and post 5 min changes in vibration thresholds in both the intervention sides- transverse and longitudinal nerve massage techniques



Graph 2 showing pre, post and post 5 min changes in heat thresholds in both the intervention sides- transverse and longitudinal nerve massage techniques



Graph 3 showing pre, post and post 5 min changes in cold thresholds in both the intervention sides- transverse and longitudinal nerve massage techniques



effects in favour of the transverse nerve massage technique which biomechanically would be produced in slider techniques and was also well tolerated by the subjects.

The following could be considered as the limitations of our study;

The study was done on asymptomatic subjects only, thus the results and findings could not be applied to symptomatic population.

Incidental sampling and also included age group could not have produced representative sample of the actual asymptomatic population.

Interventions were done on the same subject for minimizing between-subject differences of age,

gender, occupation, lifestyle and other anthropometric factors, but this could have affected external validity of our findings further.

The dosage of the intervention used in the study was not validated in earlier studies and hence another dosage or combination of techniques would have produced different results.

Both the techniques had their own beneficial effects in that they involve soft tissue mobilization together with nerve mobilization which might have some other indirect effects.

The following are the significance of our study findings;

The study is the first of its kind, especially done on the lower limb and has future implications for further studies to actually measure the mechanics of the other lower extremity nerves during nerve massage techniques in normal subjects and/or in neuropathic pain subjects using ultrasonography.⁷¹

The study was having very high internal validity due to blinding of testers about the outcomes and also observer blinding for interventions.

CONCLUSION

Tibial nerve transverse massage technique reduced the vibration threshold significantly better than the tensioner technique in asymptomatic subjects studied. Clinically these techniques also reduced the heat thresholds and increased cold thresholds in the subjects which were not statistically significant.

REFERENCES

- Butler DS. Adverse Mechanical Tension in the Nervous System: A Model for Assessment and Treatment. *Aus J Physiother.* 1989; 35: 227- 238.
- Robert J. Nee, David Butler. Management of peripheral neuropathic pain: Integrating neurobiology, neurodynamics, and clinical evidence. *Phys Ther Sport.* 2006; 7: 36-49.
- Butler DS. Mobilization of the Nervous System. Churchill Livingstone. 1996.
- T. M. Hall, R. L. Elvey. Nerve trunk pain: physical diagnosis and treatment. *Man Ther.* 1999; 4: 63- 73.
- Shacklock MO. Editorial- Improving application of neurodynamic (neural tension) testing and treatments: A message to researchers and clinicians. *Man Ther.* 2005; 10: 175-179.
- Shacklock MO. Clinical Neurodynamics- A New System of Musculoskeletal Treatment. Elsevier, Oxford. 2005.
- Coppieters MW, Butler DS. Do 'sliders' slide and 'tensioners' tension? An analysis of neurodynamic techniques and considerations regarding their application. *Manual Therapy* (2007), doi:10.1016/j.math. 2006; 12: 008.
- Topp KS, Boyd BS. Structure and biomechanics of peripheral nerves: nerve responses to physical stresses and implications for physical therapist practice. *Phys Ther.* 2006; 86: 92-109.
- Millesi H, Zoch G and Reihnsne R. Mechanical Properties of Peripheral Nerves. *Clin Orthop Rel Res.* 1995; 314: 76-83.
- Dilley A, Lynn B, Pang SJ. Pressure and stretch mechanosensitivity of peripheral nerve fibres following local inflammation of the nerve trunk. *Pain.* 2005; 117: 462-472.
- B. van der Heide, G. T. Allison, M. Zusman. Pain and muscular responses to a neural tissue provocation test in the upper limb. *Manual Therapy.* 6: 154- 162.
- M. W. Coppieters, K. H. Stappaerts, F. F. Staes, D. G. Everaert. Shoulder girdle elevation during neurodynamic testing: an assessable sign? *Manual Therapy.* 6: 88- 96
- Andrea Julius, Rebecca Lees, Andrew Dilley and Bruce Lynn. Shoulder posture and median nerve sliding. *BMC Musculoskeletal Disorders.* 5: 23.
- Wilson S, Selvarathnam PJ, and Briggs C. Strain at the Subclavian artery during the upper limb tension test. *Australian Journal of Physiotherapy.* 40: 243-248.
- SM Balster, G A Jull. Upper trapezius muscle activity during the brachial plexus tension test in asymptomatic subjects. *Manual Therapy.* 2: 144- 149.
- Coppieters MW, Alshami AM, Hodges PW. An experimental pain model to investigate the specificity of the neurodynamic test for the median nerve in the differential diagnosis of hand symptoms. *Arch Phys Med Rehabil.* 87: 1412-7.

17. Yaxley GA, and Jull GA. A modified upper limb tension test: An investigation of responses in normal subjects. *Australian Journal of Physiotherapy*. 37: 143- 152.
18. Moses AS and Carman JB. Anatomy of the cervical spine: Implication for the upper limb tension test. *Australian Journal of Physiotherapy*. 42: 31-35.
19. Michel W Coppieters, Karel H Stappaerts, Dirk G Everaert, Filip F Staes. Addition of Test Components During Neurodynamic testing: Effect on Range of Motion and Sensory Responses. *J Orthop Sports Phys Ther*. 13: 226-237.
20. P C Lew, C A Briggs. Relationship between the cervical component of the slump test and change in hamstring muscle tension. *Manual Therapy*. 2: 98-105.
21. Evan K. Johnson, Cynthia M. Chiarello. The Slump Test: The Effects of Head and Lower Extremity Position on Knee Extension. *J Orthop Sports Phys Ther*. 26: 310-317.
22. Boland RA and Adams RD. Effects of ankle dorsiflexion on range and reliability of straight leg raising. *Australian Journal of Physiotherapy*. 46: 191-200.
23. Dilley A, Lynn B, Greening J, DeLeon N. Quantitative in vivo studies of median nerve sliding in response to wrist, elbow, shoulder and neck movements. *Clin Biomech (Bristol, Avon)*. 2003; 18(10): 899-907.
24. Yaxley GA, and Jull GA. Adverse Tension in the Neural System- A Preliminary Study of Tennis Elbow. *Australian Journal of Physiotherapy*. 39: 15-22.
25. Kelly SM and Jull GA. Breast surgery and neural tissue Mechanosensitivity. *Australian Journal of Physiotherapy*. 44: 31-37.
26. Hall TM, Quintner JL. Responses to mechanical stimulation of the upper limb in painful cervical radiculopathy. *Australian Journal of Physiotherapy*. 42: 277-285.
27. Michael O Shacklock. Positive Upper Limb Tension Test in a case of surgically proven neuropathy: Analysis and Validity. *Manual Therapy*. 1: 154-161.
28. M. Sterling, J. Treleaven, G. Jull. Responses to a clinical test of mechanical provocation of nerve tissue in whiplash associated disorder. *Manual Therapy*. 7: 89-94.
29. Yeung E, Jones M and Hall B. Response to the slump test in a group of female whiplash patients. *Australian Journal of Physiotherapy*. 43: 245- 252.
30. Michel W. Coppieters, Kimberly Kurz, Thor Einar Mortensen, Nicola L. Richards, Ingrid A. Skaret, Laurie M. McLaughlin, Paul W. Hodges. The impact of neurodynamic testing on the perception of experimentally induced muscle pain. *Manual Therapy*. 10: 52-60.
31. Sharon E. Turl, Keith P. George. Adverse Neural Tension: A Factor in Repetitive Hamstring Strain? *J Orthop Sports Phys Ther*. 27: 16-21.
32. S Pahor, R Toppenberg. An investigation of neural tissue involvement in lateral ankle sprains. *Manual Therapy*. 1: 192-197.
33. Hough AD, Moore AP, Jones MP. Reduced longitudinal excursion of the median nerve in carpal tunnel syndrome. *Arch Phys Med Rehabil*. 2007; 88: 569-76.
34. Kleinrensink GJ, Stoeckart R, Mulder PG, Hoek G, Broek T, Vleeming A, Snijders CJ. Upper limb tension tests as tools in the diagnosis of nerve and plexus lesions. Anatomical and biomechanical aspects. *Clin Biomech (Bristol, Avon)*. 2000; 15 (1): 9-14.
35. Peter J. Keir, David M. Rempel. Pathomechanics of Peripheral Nerve Loading: Evidence in Carpal Tunnel Syndrome. *J Hand Ther*. 2005; 18:259-269.
36. Mahmud MA, Merlo AR, Gomes I, Becker J, Nora DB. Relationship between adverse neural tension and nerve conduction studies in patients with symptoms of the carpal tunnel syndrome. *Arq Neuropsiquiatr*. 2006; 64(2A): 277-82.
37. Jane Greening, Andrew Dilley, Bruce Lynn. In vivo study of nerve movement and mechanosensitivity of the median nerve in whiplash and non-specific arm pain patients. *Pain*. 115: 248-253.
38. J Byng. Overuse syndromes of the upper limb and the upper limb tension test: a comparison between patients, asymptomatic keyboard workers and asymptomatic non-keyboard workers. *Manual Therapy*. 2: 157-164.
39. Akalin E, ElO", PekerO", S, enocakO", Tamci S, , Gu" lbahar S, C, akmur R, O"ncel S: Treatment of carpal tunnel syndrome with nerve and tendon gliding exercises. *Am J Phys Med Rehabil*. 2002; 81:108-113.
40. Tal-Akabi A, Rushton A. An investigation to compare the effectiveness of carpal bone mobilisation and neurodynamic mobilisation as methods of treatment for carpal tunnel syndrome. *Man Ther*. 2000; 5: 214-222.

41. Michlovitz S L. Conservative Interventions for Carpal Tunnel Syndrome. *J Orthop Sports Phys Ther.* 2004; 34: 589-600.
42. Baysal O, Altay Z, Ozcan C, Ertem K, Yologlu S, Kayhan A. Comparison of three conservative treatment protocols in carpal tunnel syndrome. *Int J Clin Pract.* 2006; 60: 820-8.
43. Pinar L, Enhos A, Ada S, Gungor N. Can we use nerve gliding exercises in women with carpal tunnel syndrome? *Adv Ther.* 2005; 22: 467-75.
44. Muller M, Tsui D, Schnurr R, Biddulph-Deisroth L, Hard J. Effectiveness of Hand Therapy Interventions in Primary Management of Carpal Tunnel Syndrome: A Systematic Review. *J Hand Ther.* 2004; 17: 210-228.
45. Totten PA, Hunter JM. Therapeutic techniques to enhance nerve gliding in thoracic outlet syndrome and carpal tunnel syndrome. *Hand Clin.* 1991; 7(3): 505-20.
46. Goodyear-Smith F, Arroll B. What Can Family Physicians Offer Patients With Carpal Tunnel Syndrome Other Than Surgery? A Systematic Review of Nonsurgical Management. *Ann Fam Med.* 2004; 2: 267-273.
47. Burke F D, Ellis J, McKenna H, Bradley M J. Primary care management of carpal tunnel syndrome. *Postgrad Med J.* 2003; 79: 433-437.
48. Sweeney J, Harms A. Persistent mechanical allodynia following injury of the hand. Treatment through mobilization of the nervous system. *J Hand Ther.* 1996; 9: 328-38.
49. Kostopoulos D. Treatment of carpal tunnel syndrome: a review of the non-surgical approaches with emphasis in neural mobilization. *J Bodywork Movement Therapies.* 2004; 8: 2-8.
50. Lund A T, Amadio P C. Treatment of Cubital Tunnel Syndrome: Perspectives for the Therapist. *J Hand Ther.* 2006; 19: 170-9.
51. Coppieters M W, Bartholomeeusen K E, and Stappaerts K H. Incorporating Nerve Gliding Techniques in the Conservative Treatment of Cubital Tunnel Syndrome. *J Manipulative Physiol Ther.* 2004; 27: 560- 568.
52. Cleary C K. Management of Radial Tunnel Syndrome: A Therapist's Clinical Perspective. *J Hand Ther.* 2006; 19: 186-91.
53. Trudel D, Duley J, Zastrow I, Kerr E W, Davidson R, MacDermid J C. Rehabilitation for Patients with Lateral Epicondylitis: A Systematic Review. *J Hand Ther.* 2004; 17: 243-266.
54. Ekstrom R A, Holden R. Examination of and intervention for a patient with chronic lateral elbow pain with signs of nerve entrapment. *Phys Ther.* 2002; 82: 1077-1086.
55. Crosby C A, Wehbe M A. Conservative treatment for thoracic outlet syndrome. *Hand Clin.* 2004; 20: 43-9.
56. Wehbe M A, Schlegel J M. Nerve gliding exercises for thoracic outlet syndrome. *Hand Clin.* 2004; 20: 51-5.
57. Zvulum I. Mobilizing the nervous system in cervical cord compression. *Man Ther.* 1998; 3: 42- 47.
58. Murphy D R, Hurwitz E L, Gregory A and Clary R. A Nonsurgical Approach to the Management of Patients with Cervical Radiculopathy: A Prospective Observational Cohort Study. *J Manip Physiol Ther.* 2006; 29: 279-287.
59. Coppieters M W, Stappaerts K H, Wouters L L, and Janssens K. Aberrant protective force generation during Neural provocation testing and the effect of Treatment in patients with neurogenic Cervicobrachial pain. *J Manip Physiol Ther.* 2003; 26: 99-106
60. Elvey R L. Treatment of arm pain associated with abnormal brachial plexus tension. *Aus J Physiother.* 1986; 32: 225-230.
61. Haddick E. Management of a Patient With Shoulder Pain and Disability: A Manual Physical Therapy Approach Addressing Impairments of the Cervical Spine and Upper Limb Neural Tissue. *J Orthop Sports Phys Ther.* 2007; 37: 342-350.
62. Allison G T, Nagy B M, Hall T. A randomized clinical trial of manual therapy for cervico-brachial pain syndrome – a pilot study. *Man Ther.* 2002; 7: 95-102.
63. Cleland J A, Childs J D, Palmer J A, Eberhart S. Slump stretching in the management of non-radicular low back pain: A pilot clinical trial. *Man Ther.* 2006; 11: 279-286.
64. Koury M J, Scarpelli E. A manual therapy approach to evaluation and treatment of a patient with a chronic lumbar nerve root irritation. *Phys Ther.* 1994; 74: 548-560.
65. George S Z. Characteristics of patients with lower extremity symptoms treated with slump stretching: a case series. *J Orthop Sports Phys Ther.* 2002; 32: 391-398.

66. Scrimshaw S V, Maher C G. Randomized Controlled Trial of Neural Mobilization After Spinal Surgery. *Spine*. 2001; 26: 2647-2652.
67. Ridehalgh C, Greening J, Petty N J. Effect of straight leg raise examination and treatment on vibration thresholds in the lower limb: a pilot study in asymptomatic subjects. *Man Ther*. 2005; 10: 136-143.
68. Vibrotherm™ Instruction Manual- Diabetic Foot Kare India, Chennai. 2006.
69. Rolke R et al. Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): Standardized protocol and reference values. *Pain*. 2006; 123: 231-243.
70. T. Hall, M. Zusman, R. Elvey. Adverse mechanical tension in the nervous system? Analysis of straight leg raise. *Manual Therapy*. 1998; 3: 140-146.
71. A.D. Hough, A.P. Moore, M.P. Jones. Measuring longitudinal nerve motion using ultrasonography. *Manual Therapy*. 5: 173- 180.
72. Jabre, J.F., 1994. "Nerve rubbing" in the symptomatic treatment of ulnar nerve paresthesia. *Muscle and Nerve*. 17: 1237.