
## Non Contact IR Foot Thermometer - FT4

Inflammation is one of the earliest signs of tissue injury and ulceration. However the clinical signs of the inflammation are usually too subtle to be detected by patients or even by the trained health care providers. Skin temperature measurements can be easily performed and assessed by using a IR Thermometer as a diagnostic tool for diabetic foot ulcerations and Charcot arthropathy.

Patients with diabetic neuropathy who monitor foot skin temperature daily can reduce the risk of ulceration more than 4 fold according to Lawrence A, Lavery, D.P.M and colleagues.

Temperature are measured on the sole of the foot (first, third, and fifth metatarsal head; great toe; central mid-foot; and heel) and dorsum. Higher temperatures were considered to indicate ongoing inflammation. If the temperature of corresponding sites on the left and right foot differed by >2.2°C (4°F), patients were advised to contact the doctor and to reduce the number of steps taken in the following days until the temperature difference was <2.2°C. Early diagnosis and early treatment is crucial for the healing of diabetic foot lesions and resources for early interventions must therefore be available to take care of a higher number of suspected foot complications.





For more information contact



## Diabetik Foot Care India Pvt Limited

No.18/1, Kannappanagar, 3rd Main Road, Thiruvanmiyur, Chennai-41, INDIA. Tel - 91-44-4356 4129 / 4359 0510 / 93806 21607 E-mail: mesmedi@gmail.com; elango@mesmedi.com

URL: www.diabeticfootcareindia.com; www.diabeticneuropathy.co.in

